

Targeting diminished cerebral glucose metabolism for Alzheimer's disease


As demographics of developed countries shift toward an older population, the number of cases of Alzheimer's disease is anticipated to increase dramatically. Currently, there are no disease modifying treatments and approved drugs provide only symptomatic relief. Amyloid beta has been the primary drug target over the last several years, yet has not yielded any new drug approvals. Another feature of Alzheimer's disease, diminished cerebral glucose metabolism, may offer new targets for drug development.

By Dr Henderson

Alzheimer's disease (AD) is a devastating age-associated neurodegenerative disease characterised by a progressive decline in cognitive abilities. Clinical signs of AD include memory loss that disrupts daily life, difficulty in completing daily tasks and difficulties with language. Each of these symptoms progress until the patient becomes incapacitated. The prevailing theory to explain the pathophysiology of AD is the amyloid cascade hypothesis. This hypothesis is based on both the neuropathology and genetic links to early onset forms of the disease. The neuropathology of AD is characterised by the deposition of fibrillar amyloid- β (A β) peptides, amyloid plaques (AP) and accumulation of neurofibrillary tangles (NFT), mainly composed of tau protein. The prevalence of A β species in the brains of patients with AD has led support to the amyloid cascade hypothesis as the fundamental causative

mechanism in the pathogenesis of AD. This hypothesis has been further reinforced by the association between early onset AD and rare mutations in the genes *APP* (Amyloid Precursor Protein), *PSEN1* (Presenilin-1) and *PSEN2* (Presenilin-2). Mutations in these genes lead to early-onset (<60 years) familial AD, which represents about 5% of AD cases. A prolific research effort has demonstrated that these mutations result in altered processing of the APP protein and generation of toxic forms of A β . With this combination of genetics and pathology, the amyloid hypothesis has remained the dominant theory in the field and has attracted the lion's share of drug development¹.

Unfortunately, the amyloid cascade hypothesis has yet to yield any treatments for AD. Drugs candidates have targeted various aspects of A β generation and clearance, ranging from gamma secretase inhibitors designed to limit the production of toxic



forms of A β and more recently, immunotherapy approaches designed to specifically remove A β from the brain. These studies have uniformly failed to achieve primary endpoints in Phase III trials. While the amyloid cascade hypothesis has solid footing in both genetics and pathology of the disease, and considerable resources are still being devoted to it, it may be time to seriously consider other aspects of AD pathology that may offer another angle to tackle this difficult disease.

One such aspect is diminished cerebral glucose metabolism (DCGM). DCGM is a prominent, well-characterised feature of AD². Starting in the 1980s, studies examining brain glucose metabolism in patients with AD by FDG-PET found DCGM in patients with probable AD compared to normal controls (Figure 1). In these studies, significant correlations were found between DCGM and worsening performance on measures of cognitive function. The DCGM found in AD is not a general decrease in glucose utilisation but instead is associated with specific regions of the brain. Most notably, DCGM is found in the posterior cingulate and parietal, temporal and prefrontal cortices. This pattern of DCGM has been identified in at-risk individuals well before clinical signs of dementia become evident, and even in young adults. Interestingly, the regions afflicted with DCGM are associated with the brain's default network, which is a series of connected regions of the brain that are most active when one is resting quietly with closed eyes. The default network is thought to function in spontaneous cognition; thoughts that do not require outside stimuli. These activities include planning for the future, daydreaming and importantly, memory retrieval. Various studies have mapped the default network to an interconnected network comprising the prefrontal cortex, posterior cingulate,

parietal lobule and temporal cortex. Several studies have suggested that the default network may be predisposed to develop AD pathology due to the high rates of glucose metabolism found compared to other brain regions³.

The characteristic pattern of DCGM is an early event in AD and has been noted in both mild cognitive impairment (MCI) and pre-clinical AD. Mild cognitive impairment is characterised by subjective complaints of memory problems, which are not serious enough to interfere with daily living and is considered a risk state for later development of dementia. As many as 50% of MCI patients will progress to dementia, particularly AD. The potential relationship between DCGM to AD can be demonstrated in longitudinal studies which demonstrate the progressive worsening of DCGM as individuals progress from pre-symptomatic AD to MCI and ultimately to AD.

Despite the early occurrence of DCGM, the precise cause remains elusive. Historically, the DCGM seen in AD has been viewed as representing loss of synaptic activity either through cell atrophy or reduction in synaptic and dendritic fields. However, increasing evidence has pointed to a fundamental cellular or metabolic defect in AD, and perhaps a primary role of DCGM in the etiology of the disease. Several recent studies in transgenic mouse models of AD, suggest that metabolic disturbances occur before the widespread generation or deposition of A β and may be related to deficits in cellular trafficking of mitochondria. These findings are consistent with the roles of *APP* and *PSEN* in intracellular trafficking. The *APP* and *PSEN* protein complexes are thought to play a role in facilitating the movement of lipid rich cargos up and down axons. Disruption of this trafficking within the narrow confines of an axon has been proposed to result in 'neuronal traffic jams' that

Figure 1
Diminished Cerebral Glucose Metabolism (DCGM) in Alzheimer's disease. Differences between Alzheimer's disease patients and age-matched controls in brain glucose utilisation (FDG-PET, red) and atrophy (MRI, blue). Areas of overlap are shown in violet. Adapted from Frisch et al 2013²

Ketogenic diets

Ketogenic diets (KDs) became of interest in the 1920s when they were developed to reduce the occurrence of seizures in epileptics. The rationale for this type of diet was based on the ancient observation that fasting reduced seizures. In the 5th century, Hippocrates noted that a man suffering from epilepsy was cured by abstaining from all food and drink. In the King James Version of the Bible, a story describes how Jesus cured an epileptic boy (Figure 3). When asked how he did it, Jesus responded that, “this kind can come out by nothing but prayer and fasting”. KDs were developed to mimic the physiological changes seen in extended fasting. KDs are very low in carbohydrates and proteins and very high in fat and have been used successfully for many years to treat refractive childhood epilepsy.

Ketogenic diets have been employed in several neurological disorders including mild cognitive impairment. In 2010, Krikorian and co-authors reported the results of a study of 23 older adults with memory complaints placed on a KD for six weeks. After six weeks on the KD memory performance improved and the improvement correlated with the patient’s level of ketosis⁵. Unfortunately, the strict limitation on carbohydrate intake makes adherence to a KD challenging. This is particularly true for patients with AD who have documented shifts in food preference toward sweet foods, making compliance to a KD difficult.

Ketogenic agents

An alternative to a KD is to induce ketosis without change in diet. One way this has been done is with special fats called medium chain triglycerides (MCTs). MCTs are triglycerides comprised of fatty acid chains between 5-12 carbons. Fatty acids normally encountered in the diet are long chain fatty acids (LCT) with chains between 18-22 carbons. Due to the short chain lengths of medium chain fatty acids, MCTs are not subject to the regulation imposed on long chain fatty acids and are well known for their ketogenic properties. Importantly, the oxidation of medium chain fatty acids occurs regardless of carbohydrate in the diet and can induce ketosis without modification of the diet.

Two studies have been reported using MCTs to induce ketosis in mild to moderate AD patients. The first study was a single dose, crossover design. The second was a 90-day study with once-a-day dosing. Each of these studies demonstrated improvement in cognition in a predefined genetic sub population of the participants⁶. Such studies are encouraging and offer the possibility that the



Figure 3

Jesus cures an epileptic boy. Engraving by Otto Elliger depicts Jesus healing an epileptic boy by casting out a demon, after his disciples were unable to do so. Courtesy of the Digital Image Archive, Pitts Theology Library, Candler School of Theology, Emory University

induction of ketosis may prove a relatively low risk intervention in mild to moderate AD. Larger more comprehensive studies are currently under way with this approach.

Insulin signalling

Another target is the insulin and insulin-like growth factor signalling pathways. As discussed above, several lines of evidence suggest that AD is fundamentally a metabolic disease in which brain glucose utilisation and energy production are impaired. Several lines of evidence suggest that the metabolic disturbance is due to disruption in brain insulin and insulin-like growth factor (IGF) signalling pathways. Such pathways regulate neuronal survival and energy production. The development of resistance in these pathways, either through dietary factors or activities of A β , could lead to progressive disruption in energy generation and cell survival. This hypothesis is sometimes referred to as type 3 diabetes or ‘Brain Diabetes’⁷.

Studies done targeting insulin and insulin-like signalling have met with some success. Notable among these interventions is the application of nasal insulin. As with diabetes, the resistance to insulin can be overcome by provision of more insulin. Systemic provision of insulin would have profound effects on circulating glucose; therefore, to avoid possible harmful side-effects, insulin has been administered intranasal to target the brain directly. Published studies have found that such treatments increase brain insulin levels and improve performance memory tasks while having little effect on plasma glucose and insulin levels.

References

- 1 Holtzman, DM, Mandelkow, E and Selkoe, DJ. Alzheimer disease in 2020. Cold Spring Harb Perspect Med, 2012. 2(11).
- 2 Frisch, S et al. Dissociating memory networks in early Alzheimer's disease and frontotemporal lobar degeneration – a combined study of hypometabolism and atrophy. PLoS One, 2013. 8(2): p. e55251.
- 3 Buckner, RL et al. Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. J Neurosci, 2005. 25(34): p. 7709-17.
- 4 Trushina, E et al. Defects in mitochondrial dynamics and metabolomic signatures of evolving energetic stress in mouse models of familial Alzheimer's disease. PLoS One, 2012. 7(2): p. e32737.
- 5 Krikorian, R et al. Dietary ketosis enhances memory in mild cognitive impairment. Neurobiol Aging, 2010. Online.
- 6 Henderson, ST et al. Study of the ketogenic agent AC-1202 in mild to moderate Alzheimer's disease: a randomized, double-blind, placebo-controlled, multicenter trial. Nutr Metab (Lond), 2009. 6(1): p. 31.
- 7 de la Monte, SM. Brain insulin resistance and deficiency as therapeutic targets in Alzheimer's disease. Curr Alzheimer Res, 2012. 9(1): p. 35-66.
- 8 Craft, S et al. Intranasal Insulin Therapy for Alzheimer Disease and Amnesic Mild Cognitive Impairment: A Pilot Clinical Trial. Arch Neurol, 2011.

Larger, more comprehensive studies are currently under way with this approach⁸.

Conclusions

With only the currently available treatments, the prevalence of AD is predicted to increase dramatically in the next 40 years. Current treatments, while effective, do not significantly slow the progression of the disease, and new therapies are desperately needed. A tremendous research effort over the last 10 years has greatly increased our understanding of AD. Many exciting therapies have been developed over the years, but have not produced the results we would all like to see in the clinic.

Addressing another aspect of AD pathology, such as DCGM, is a relatively new area of research yet offers the promise of many different avenues of attack. Addressing low glucose availability by the induction of ketosis is one such promising approach. Addressing the insulin/IGF signalling pathway is another. Fortunately, both of these approaches have entered larger clinical trials and results are expected within a few years. **DDW**

Trained in Genetics, Dr Henderson is now Vice President of Research and Development at Accera, a biotechnology company focused on cognition and the use of ketone bodies in neurodegenerative diseases. He received his B.A. in Biology from Washington University, St. Louis, MO and his Ph.D. in Molecular Genetics and Cell Biology from University of Chicago. He received extensive post-doctoral training in developmental biology at the University of Wisconsin. Dr. Henderson is the scientific founder of Accera.